

Early Exposure to Common Anesthetic Agents Causes Widespread Neurodegeneration in the Developing Rat Brain and Persistent Learning Deficits

Vesna Jevtovic-Todorovic,¹ Richard E. Hartman,² Yukitoshi Izumi,³ Nicholas D. Benschoff,³ Krikor Dikranian,³ Charles F. Zorumski,³ John W. Olney,³ and David F. Wozniak³

¹Department of Anesthesiology, University of Virginia Health System, Charlottesville, Virginia 22908, and Departments of ²Neurology and ³Psychiatry, Washington University School of Medicine, St. Louis, Missouri 63110

Recently it was demonstrated that exposure of the developing brain during the period of synaptogenesis to drugs that block NMDA glutamate receptors or drugs that potentiate GABA_A receptors can trigger widespread apoptotic neurodegeneration. All currently used general anesthetic agents have either NMDA receptor-blocking or GABA_A receptor-enhancing properties. To induce or maintain a surgical plane of anesthesia, it is common practice in pediatric or obstetrical medicine to use agents from these two classes in combination. Therefore, the question arises whether this practice entails significant risk of inducing apoptotic neurodegeneration in the developing human brain. To begin to address this problem, we have administered to 7-d-old infant rats a combination of drugs commonly used in pediatric anesthesia (midazolam, nitrous oxide, and isoflurane) in doses sufficient to maintain a surgical plane of anesthesia for 6 hr, and have observed that this causes widespread apoptotic neurodegeneration in the developing brain, deficits in hippocampal synaptic function, and persistent memory/learning impairments.

Key words: NMDA antagonists; GABA agonists; isoflurane; midazolam; nitrous oxide; apoptosis

Introduction

Advances in pediatric and obstetric surgery have resulted in an increased complexity, duration, and number of anesthesia procedures. To minimize risks, it is necessary to understand the effects of anesthetic drugs on the developing nervous system. Presently used anesthetics act by two principal mechanisms: (1) an increase in inhibition via GABA_A receptors (e.g., benzodiazepines, barbiturates, propofol, etomidate, isoflurane, enflurane, and halothane) (Franks and Lieb, 1994), and (2) a decrease in excitation through NMDA receptors [e.g., ketamine, nitrous oxide (N₂O), and xenon] (Lodge and Anis, 1982; Franks et al., 1998; Jevtovic-Todorovic et al., 1998; Mennerick et al., 1998). Recent findings indicate that drugs that act by either of these mechanisms induce widespread neuronal apoptosis in immature rat brain when administered during synaptogenesis (Ikonomidou et al., 1999, 2000; Ishimaru et al., 1999). In addition, ethanol, an agent with both NMDA antagonist and GABA_{mimetic} properties, triggers a widespread pattern of apoptotic neurodegeneration in the developing rat brain, which is a composite of the patterns induced by NMDA antagonist and GABA_{mimetic} drugs (Ikonomidou et al., 1999). These findings raise questions regarding the potential risk posed by currently used anesthesia protocols. To address this issue, we studied the histopathological, electrophysiological, and behavioral effects of exposure of 7-d-old

rats to N₂O, isoflurane, and midazolam, three agents commonly used in obstetric and pediatric anesthesia.

Materials and Methods

Animals. Seven-day-old male and female Sprague Dawley rats were used for all experiments. At postnatal day 7 (P7), experimental rats were exposed to 6 hr of anesthesia, and controls were exposed to 6 hr of mock anesthesia. They were then allowed to recover and were divided into three randomly selected groups. One group was used for histopathology studies at several acute postanesthesia intervals. The second group was used for behavioral studies, which involved evaluating the rats on several tests over a 160 d period. The third group was used to study long-term potentiation (LTP) in hippocampal slices at P29–P33. Daily inspection and weighing was performed on the latter two groups at P6–P21 to evaluate general health and development.

Anesthesia treatment. N₂O and oxygen were delivered using a calibrated flowmeter. Midazolam was dissolved in 10% DMSO immediately before administration. For control experiments, 10% DMSO was used. To administer specific concentrations of N₂O, both normobaric [1 atmosphere (atm), 100 vol%] and hyperbaric conditions were required. In experiments requiring hyperbaric conditions, the N₂O/oxygen mixture was introduced at a pressure of 2.0 atm (200 vol% = 150 vol% N₂O and 50 vol% oxygen) and sustained for the duration of the experiment (Mahmoudi et al., 1989; Gonsowski and Eger, 1994). A relief valve on the hyperbaric chamber allowed continuous escape of gases to avoid accumulation of carbon dioxide. Animals were kept normothermic throughout experiments. For N₂O concentrations of <80 vol%, normobaric conditions were used. The N₂O/oxygen mixture was delivered to the chamber while a relief valve was kept open so that pressure inside the chamber remained at 1 atm. For control experiments, air was substituted for the gas mixtures. For experiments with isoflurane, we used an agent-specific vaporizer that delivers a set percentage of anesthetic into the chamber. After initial equilibration of the N₂O/oxygen/isoflurane or air/isoflurane atmosphere inside the chamber, the composition of the cham-

Received Aug. 20, 2002; revised Nov. 4, 2002; accepted Nov. 8, 2002.

This work was supported in part by National Institutes of Health Grants AG 11355, DA 05072, HD 37100, AG 18434, MH 45493, and AA 12951 and Career Development Award K08 DA 00406 (V.J.-T.). We thank Adam Myenberg, Anna Pieper, and Anthony Nardi for technical assistance.

Correspondence should be addressed to Vesna Jevtovic-Todorovic, Department of Anesthesiology, University of Virginia Health System, P.O. Box 800710, Charlottesville, VA 22908. E-mail: vj3w@virginia.edu.

Copyright © 2003 Society for Neuroscience 0270-6474/03/230876-07\$15.00/0

ber gas was analyzed by mass spectrometry for N₂O or nitrogen, isoflurane, carbon dioxide, and oxygen concentration. All experiments were approved by The Animal Use and Care Committees of the University of Virginia and Washington University School of Medicine.

Arterial blood gas analysis. To determine adequacy of ventilation, arterial blood was sampled at the end of anesthesia by obtaining a single sample (100 μ l) from the left cardiac ventricle using a 32 gauge hypodermic needle. Bicarbonate concentration (millimoles per liter), oxygen saturation (%), pH, paCO₂ (partial pressure of carbon dioxide in mmHg), and paO₂ (partial pressure of oxygen in mmHg) were measured immediately after blood collection, using a Nova Biomedical blood gas apparatus. Control samples were obtained from air/DMSO-treated pups.

Histopathological studies. All pups were deeply anesthetized and perfused with aldehyde fixatives for histopathology studies of the brains. We have found that activated caspase-3 [immunocytochemical (ICC)] is an excellent method for marking neurons that are in an early stage of apoptosis (Olney et al., 2002a,b), and DeOlmos silver staining is very useful for mapping patterns of cell death in the developing brain (Ikonomidou et al., 1999, 2000). To confirm the apoptotic nature of the cell-death process, electron microscopy is the most reliable available means (Dikranian et al., 2001). Pups used for studying caspase-3 activation or for silver staining were perfused with a mixture of paraformaldehyde (4%) in Tris buffer, pH 7.4, either 2 hr (caspase) or 18 hr (silver) after cessation of anesthesia, whereas those used for electron microscopy studies were perfused with a mixture of paraformaldehyde (4%) and glutaraldehyde (1.5%) in cacodylate buffer, pH 7.4, at 10 hr after anesthesia.

For activated caspase-3 ICC, 50- μ m-thick vibratome sections were processed by procedures described recently (Olney et al., 2002b), using a primary anti-active caspase-3 antiserum raised in rabbits (D175; Cell Signaling Technology, Beverly, MA). For silver staining, the fixed brains were cut by vibratome into 50 μ m sections and stained by the DeOlmos cupric silver method, as described previously (DeOlmos and Ingram, 1971; Corso et al., 1997). For electron microscopy, ultrathin sections were stained with uranyl acetate/lead citrate and viewed in a 100C Jeol (Peabody, MA) electron microscope.

To determine the degree of neurodegeneration in a given brain region, we used the optical dissector and fractionator method (West, 1999). A counting frame (0.05 \times 0.05 mm; dissector height, 0.05 mm) and a high numerical aperture objective lens were used to visualize neurons. Unbiased sampling of each brain region was performed by randomly selecting 10–12 viewing fields over which the counting frame was positioned for counting at different focal levels. The numerical density of degenerating neurons in any given region was determined by counting argyrophilic profiles in 50- μ m-thick sections stained by the DeOlmos silver method. Differences between anesthetic-treated and control values for a given brain region were evaluated statistically by Student's *t* test. To obtain an estimate of the severity of induced degeneration in the anesthetic-treated brains, the numerical density values for the anesthetic-treated animals were divided by those for the controls. The result was expressed as a "fold" increase over baseline, baseline being the rate at which neuronal cell death was occurring as a natural phenomenon in control brains. Counting was done by an experienced histopathologist who was unaware of the treatment condition.

Electrophysiological studies. Rats exposed for 6 hr to anesthetic agents at P7 were killed at P29–P33 for preparation of hippocampal slices using standard methods (Zorumski et al., 1996). At the time of study, slices were placed in a submersion recording chamber at 30°C. Extracellular recordings were obtained from the apical dendritic region for analysis of population EPSPs using 2 M NaCl electrodes. Evoked responses were elicited with 0.2 msec constant-current pulses through a bipolar electrode in the Schaffer collateral pathway (CA1 hippocampal field) every 30–60 sec at an intensity sufficient to elicit 50% maximal EPSPs. After establishing a stable baseline, LTP was induced by applying a single 100 Hz \times 1 sec stimulus train using the same amplitude current.

Behavioral studies. Rats exposed to the anesthetic "triple cocktail" or DMSO vehicle on P7 were evaluated behaviorally at subsequent ages using the following measures [similar to those described in greater detail by Wozniak et al. (1989, 1990, 1991), Ho et al. (2000), and Hartman et al. (2001)]: (1) Ascent test at P10, P12, and P14 (to assess the acute response to

drug treatment); (2) auditory/tactile startle and prepulse inhibition (PPI) of startle at P20; (3) a 1 hr locomotor activity test in a home-cage environment at P21; and (4) a sensorimotor battery of tests at P22, including walking initiation, ledge walking, inclined plane performance, and elevated platform performance. (5) Spatial reference memory was evaluated using the Morris water navigation test which included cued (P28), place (P32 and P131), and probe trials. (6) Spatial working memory was assessed at P53, using a win-shift spatial discrimination protocol in the radial arm maze.

Water navigation testing was conducted as two separate studies, each involving different groups of experimental and control rats ($n = 9–11$ per group). Except as is explained below, the same conditions were used for both studies, as follows: rats were tested in a small pool (100 cm inner diameter) as juveniles and in a larger pool (180 cm inner diameter) as adults. In the cued trials, rats were tested for their ability to swim to a visible platform that was switched to a new location for each trial. The place condition involved testing the rats' ability to learn the location of a platform (submerged, not visible) which remained in the same location for all trials. When subjected to place testing as juveniles (P32), the rats were given two blocks of trials (two trials per block) each day for 5 d, with an intertrial interval of 30 sec. When the rats were retested as adults (P131), using a new platform location, a protocol involving one block of trials (two trials per block) each day for 5 d was used in an effort to increase task difficulty and improve test sensitivity. Study 2 was conducted exactly like study 1, except that after 5 d of place testing as adults, rats in study 2 were tested for an additional 5 d to determine whether they could progressively improve their performance to an asymptotic level, to provide evidence of learning. Probe trials were conducted after the last place trial of juvenile testing and after the last place trials of blocks 5 and 10 during adult testing. Probe trials involved removing the platform and evaluating the rats' search patterns for the missing platform by quantifying time spent in the target quadrant and number of platform crossings.

The rats in both study 1 and study 2 were tested in the radial arm maze at P53 according to an identical win-shift spatial discrimination (working memory) protocol in each study.

Data analyses. The data from each behavioral test in both study 1 and study 2 were subjected to an ANOVA involving treatment and study as between-subjects variables. In no case was a treatment by study interaction found; therefore, for additional analysis, the data from studies 1 and 2 were combined (and the variable study was deleted), except for data pertaining to the last 5 d of place testing as adults, which were generated only in study 2. Data from the preweaning tests were typically analyzed using ANOVA models that involved treatment and litter as between-subjects variables and test sessions as a within-subjects variable. Data from postweaning tests were typically analyzed using ANOVA models with treatment and gender as between-subjects variables and blocks of trials as a within-subjects variable. Pairwise comparisons were conducted after significant effects of treatment or significant interactions involving treatment and other relevant variables, and *p* values exceeding Bonferroni corrected levels were noted when appropriate by the symbol †. The Huynh–Feldt statistic was used to adjust *p* values to help protect against violations of compound symmetry when more than two levels of a within-subjects variable were used in an ANOVA model.

Results

Neonatal anesthetic treatment does not induce metabolic or respiratory distress

To assess the effects of selected anesthetics on the developing brain, we exposed P7 rats to midazolam, isoflurane, and N₂O, either individually or in clinically relevant combinations, for 6 hr. In animals

Table 1. Arterial blood gas analysis

Arterial blood gas	Controls ($n = 8$)	Treated ($n = 8$)
pH	7.52 \pm 0.05	7.49 \pm 0.02
paCO ₂ (mmHg)	30 \pm 8	29 \pm 2.7
paO ₂ (mmHg)	117 \pm 13	93 \pm 11
HCO ₃ ⁻ (mmol/l)	23 \pm 3	22 \pm 1.5
SaO ₂ (%)	99 \pm 0.3	96 \pm 1.1

exposed for 6 hr to these drugs, either individually or in combination, including the triple cocktail (isoflurane plus N₂O plus midazolam) (Table 1), there were no signs of metabolic or respiratory distress. Oxygen saturation, paO₂, paCO₂, and pH did not differ significantly from control animals exposed to air plus vehicle for 6 hr. A slight increase in pH in both groups is attributable to transient hyperventilation caused by cardiac puncture.

Neonatal anesthetic treatment induces extensive apoptotic neurodegeneration

In vehicle-treated animals (10% DMSO plus air), both cupric silver and activated caspase-3 ICC staining revealed a sparsely scattered pattern of baseline physiological cell death (Fig. 1*a,f,h,j*). Animals treated with N₂O alone (50, 75, or 150 vol%) or midazolam alone (3, 6, or 9 mg/kg, i.p.) showed no significant increase in apoptotic neurodegeneration compared with control animals (data not shown). However, animals that were treated with isoflurane alone (0.75, 1.0, or 1.5 vol%) exhibited dose-dependent neurodegeneration ($p < 0.05$). The most vulnerable brain regions were the laterodorsal and anteroventral thalamic nuclei, where even the lowest isoflurane concentration (0.75%) caused a significant increase in neuronal degeneration (16-fold and ninefold increase, respectively). The parietal cortex (layer II) was also affected in an apparently dose-dependent manner, although the damage was significantly greater than controls at only the highest isoflurane concentration (1.5 vol%) ($p < 0.05$).

When a nontoxic dose of midazolam (9 mg/kg, i.p.) was followed by 6 hr of a low/minimally toxic concentration of isoflurane (0.75 vol%), there was a significant increase in apoptotic neurodegeneration compared with this concentration of isoflurane alone. Under this condition (double GABA-mimetic cocktail), the damage was evident primarily in the laterodorsal and anterodorsal thalamus and parietal cortex (layer II). However, when 6 hr of N₂O at a low/nontoxic concentration (75 vol%) was added to the above double cocktail, this triple cocktail (midazolam plus 6 hr of N₂O plus isoflurane) resulted in a robust neurodegenerative reaction entailing more severe damage in the thalamus and parietal cortex than was caused by the double cocktail, and also producing moderate to severe damage in many other brain regions. It warrants emphasis that all triple cocktail-exposed pups invariably sustained brain damage, and that the pattern of damage was identical in each of these brains. We have compiled a comprehensive list of the damaged brain regions (Table 2), and for each region the severity of neurodegeneration in anesthesia-exposed brains is rated in terms of how many fold greater it is compared with the baseline rate in the same region of control




Figure 1. Triple anesthetic cocktail induces apoptotic neurodegeneration. *a–l* are light micrographic scenes from various brain regions of either a control rat (*a, f, h, j*) or a rat exposed to the triple anesthetic cocktail (*b–e, g, i, k, l*). Some sections are stained by the DeOlmos silver method (*a, b, d, f, g, k*), and the remainder are immunocytochemically stained to reveal caspase-3 activation (*c, e, h–j, l*). The regions illustrated are the posterior cingulate/retrosplenial cortex (*a–c*), subiculum (*d, e*), anterior thalamus (*f, g*), rostral CA1 hippocampus (*h, i*), and parietal cortex (*j–l*). The individual nuclei shown in the anterior thalamus (*f, g*) are laterodorsal (LD), anterodorsal (AD), anteroventral (AV), anteromedial (AM), and nucleus reuniens (NR). *m* and *n* are electron micrographic scenes depicting the ultrastructural appearance of neurons undergoing apoptosis. The cell in *m* displays a very early stage of apoptosis in which dense spherical chromatin balls are forming in the nucleus at a time when the nuclear membrane remains intact and very few changes are evident in the cytoplasm. The cell in *n* exhibits a much later stage of apoptosis in which the entire cell is condensed, the nuclear membrane is absent, and there is intermixing of nuclear and cytoplasmic constituents. These are hallmark characteristics of neuronal apoptosis as it occurs in the *in vivo* mammalian brain.

brain (the baseline rate being that attributable to physiological cell death that occurs naturally in the developing brain). This list (Table 2) includes only those brain regions, or subfields within specific regions, where the rate of degeneration was at least 15-fold greater than the control rate. In all of these regions, the numerical density of degenerating neurons in the anesthetic cocktail rats was significantly greater than the control value at a level of $p < 0.001$.

The two histological methods used (activated caspase-3 ICC and silver staining) to evaluate the neurodegenerative reaction in experimental brains compared with controls were mutually confirmatory regarding the pattern of neurodegeneration induced by the triple cocktail (Fig. 1*a–l*). Electron microscopic evaluation

Table 2. Brain regions in which anesthesia-induced neurodegeneration was most heavily concentrated

Brain region	Severity (fold increase)
Medial septal nucleus	27
Diagonal band of Broca	29
Nucleus accumbens	33
Rostral caudate nucleus	26
Globus pallidus	43
Amygdaloid nuclei	
Basolateral	22
Medial	25
Cortical	23
Thalamic nuclei	
Paraventricular	34
Anterodorsal	28
Anteroventral	57
Anteromedial	51
Laterodorsal	68
Reuniens	34
Parafascicularis	45
Hippocampus, rostral CA1	21
Subiculum	58
Cingulate cortex	32
Retrosplenial cortex	53
Neocortex (layers II and IV)	
Frontal	18
Parietal	34
Temporal	22
Occipital	35
Hypothalamus	
Anterior	22
Ventromedial	34
Dorsomedial	21
Mammillary complex	33

Severity is expressed as the fold increase (i.e., how many times greater the density of degenerating neurons was in the experimental brain compared with the baseline rate of degeneration in the same region of control brain).

confirmed that the degenerating neurons displayed classical morphological changes characteristic of apoptosis (Fig. 1*m,n*).

Triple anesthetic cocktail induces profound LTP suppression

To assess long-term electrophysiological effects of early anesthetic exposure, we examined synaptic function and LTP in hippocampal slices prepared at P29–P33 from rats treated at P7 with 10% DMSO (controls), N₂O, isoflurane, midazolam, or a combination of the three agents. Baseline EPSPs elicited by single stimuli did not differ among the five groups (data not shown). LTP induction was unchanged in DMSO-treated controls compared with naive controls [changes in EPSP slope 60 min after tetanus in naive and DMSO animals were $144.1 \pm 7.9\%$ of baseline ($n = 6$) and $139.4 \pm 6.3\%$ of baseline ($n = 10$), respectively]. A less robust LTP response was observed in rats treated with N₂O ($129.1 \pm 5.9\%$; $n = 8$), isoflurane ($125.1 \pm 3.1\%$; $n = 6$), or midazolam ($120.9 \pm 2.2\%$; $n = 8$), but only midazolam-treated animals differed significantly from controls ($p = 0.022$ by *t* test). In contrast, slices from rats treated with the triple anesthetic cocktail exhibited profound suppression of LTP ($105.0 \pm 6.5\%$ of baseline; $n = 10$; $p = 0.001$), despite the presence of robust short-term potentiation (Fig. 2).

Triple anesthetic cocktail induces long-term impairment in spatial learning/memory

Because the anesthetic triple cocktail produced the most profound neurodegenerative effects, we decided to determine whether developmental exposure to this treatment also had last-




Figure 2. Effects of anesthetic exposure on LTP in the CA1 region of rat hippocampal slices. *a*, The graph depicts the time course of change in field EPSP slope (\pm SEM) in hippocampal slices from control rats treated with DMSO (\bullet) and rats exposed to the triple anesthetic combination (*triple c.*, \circ). A single 100 Hz \times 1 sec tetanus was delivered at time 0 (arrow). *b*, Traces to the right of the graph show examples of EPSPs before (solid traces) and 60 min after (dashed traces) the tetanus in slices from animals treated with the various anesthetics. Hippocampal slices were prepared at P29–P33 from rats treated at P7.

ing effects on behavior. To accomplish this, we assessed sensorimotor and cognitive function from early stages of development into adulthood and found that the anesthetic cocktail rats did not differ from controls in (1) overall growth as indexed by daily body weight measurements, (2) sensorimotor ability as measured by the ascent test (preweaning) and sensorimotor battery (postweaning), (3) acoustic or tactile startle reactivity or PPI as measured by startle/PPI testing, or (4) various movement-related variables, as measured during a 1 hr locomotor activity test.

When tested in the Morris water maze on P28, anesthetic cocktail and control rats performed similarly in terms of path length and latency when learning to swim to a visible platform during cued trials (data not shown). However, when we examined spatial reference memory capabilities in the same animals during place trials (submerged platform, fixed location) on P32, anesthetic cocktail rats showed significant acquisition deficits (Fig. 3*a*). The deficit was evidenced by the anesthetic cocktail rats showing slower acquisition rates during the middle blocks of trials, although they improved and performed like controls by the end of training and also exhibited control-like performance levels during probe trials (data not shown). When retested on the water maze as adults (P131), using a more difficult place-learning protocol, anesthetic cocktail rats again demonstrated impaired performance during place trials (Fig. 3*b*) but, unlike the earlier findings, they also demonstrated impaired retention performance during probe trials. Specifically, they spent significantly less time searching the pool quadrant where the platform had been (time in target quadrant) (Fig. 3*c*) and crossed over the former platform location significantly fewer times than controls (data not shown). Moreover, the control rats from study 2 that were subjected to an additional 5 d of testing improved their performance to asymptotic levels, suggesting that learning occurred, whereas the anesthetic cocktail group from study 2 showed no such improvement (Fig. 3*b*). In addition, the control rats showed significantly higher levels of retention during the probe trials in terms of target quadrant time (Fig. 3*c*) and platform crossings (data not shown). Swimming speeds were also analyzed during cued and place trials, and no differences were observed, further suggesting that swimming performance deficits were not responsible for the place-learning impairments in the anesthetic cocktail rats (data not shown).

When spatial working memory capabilities were tested in the radial arm maze at P53, anesthetic cocktail rats were significantly

ory as adults. The lack of differences between groups during the cued trials or with regard to swimming speeds as well as the absence of performance differences on a wide array of behavioral tests suggest that the learning/memory deficits in the anesthetic cocktail rats did not result from nonassociative performance factors such as sensorimotor disturbances, altered emotionality, or changes in motivation. Our additional finding that the anesthesia-exposed rats had lasting deficits in hippocampal synaptic function may help explain the learning deficits, in that the hippocampus is well known to play an important role in memory and learning. However, it would probably be simplistic to attribute these deficits solely to hippocampal damage, in view of substantial evidence (Aggleton and Brown, 1999; Mitchell et al., 2002) that fundamental memory functions are not mediated by the hippocampus alone, but rather by a distributed network that includes, in addition to the hippocampus, anterior thalamic nuclei, mammillary bodies, and retrosplenial cortex. Each of the latter three structures was damaged in the anesthesia-exposed brains more severely than the hippocampus.

The triple anesthetic drug protocol used in these experiments is one that is commonly used in pediatric anesthesia practice. We selected a dose of midazolam that is considered sedating (sedative dose is between 1 and 10 mg/kg, i.p., in rats) (Kissin et al., 1990) and followed this with only those concentrations of N₂O and isoflurane that are required to induce and maintain a surgical plane of anesthesia. The minimum alveolar anesthetic concentration that prevents purposeful movement to supramaximal noxious stimulation in 50% of subjects (MAC) for either N₂O or isoflurane in humans is ~67% of the MAC in rats (Hornbein et al., 1982; Mahmoudi et al., 1989; Stevens and Kingston, 1992; Orliaguet et al., 2001). Thus, the concentrations of N₂O (75%) and isoflurane (0.75%) used in our rat study would be comparable with exposing a human to 50% N₂O and 0.5% isoflurane, concentrations that are well within clinically used ranges.

It has been shown in previous studies (Ikonomidou et al., 1999, 2000) that peak vulnerability to the apoptogenic action of NMDA antagonists and GABA mimetics is during the synaptogenesis period, also known as the brain growth-spurt period. This is a readily recognized period during which the brain grows at an accelerated rate because newly differentiated neurons throughout the brain are rapidly expanding their dendritic arbors to provide the required surface area to accommodate new synaptic connections. The brain growth spurt occurs in different mammalian species at different times relative to birth. In rats, it begins a day or two before birth and ends ~2 weeks after birth, whereas in humans it starts at the beginning of the third trimester and ends several years after birth (Dobbing and Sands, 1979). Comparing the brain growth curves for rats and humans, the period of peak brain growth occurs in rats between the fourth and tenth postnatal days and in humans between the last month of gestation and first 6 months after birth. Although neurodevelopmental age equivalencies between rats and humans cannot be specified with precision, our decision to use 7-d-old rats for the present study was based on the assumption that this neurodevelopmental age in the rat is equivalent to the human age from 0 to 6 months after a term birth, or perhaps from 0 to ≥12 months after a premature birth. It is quite common, often out of necessity, but sometimes on an elective basis, for surgical procedures requiring general anesthesia to be performed on human infants in this neurodevelopmental age range.

Various anesthesia protocols have been used in pediatric medicine for many decades without clear evidence linking anesthesia

exposure to subsequent neurobehavioral disturbances. Establishing such a link is rendered difficult by many confounding variables. For example, human infants who undergo general anesthesia often have a history of prematurity and/or adverse peripartum events, including prolonged exposure to sedatives or anticonvulsants in an intensive care unit. Moreover, linking neurobehavioral disturbances to perinatal drug exposure is difficult unless gross signs of dysmorphogenesis are present. For example, researchers were first alerted to the fetal alcohol syndrome by conspicuous dysmorphogenic effects (craniofacial malformations), and this led to the subsequent recognition that fetal exposure to ethanol can cause a wide range of neurobehavioral disturbances in the absence of dysmorphogenic effects (Streissguth and O'Malley, 2000). Our findings indicate that an anesthesia protocol that does not alter somatic development or induce sensorimotor impairments in rats does result in learning/memory deficits that are subtle enough to be easily overlooked. Additional animal research will be needed to address the potential developmental neurotoxicity of other clinically used anesthetic protocols. In addition, well designed clinical investigations are needed to assess the potential relevance of the animal findings to obstetric or pediatric anesthesia.

References

- Aggleton JP, Brown MW (1999) Episodic memory, amnesia, and the hippocampal-anterior thalamic axis. *Behav Brain Sci* 22:425–489.
- Corso TD, Sesma MA, Tenkova TI, Der TC, Wozniak DF, Farber NB, Olney JW (1997) Multifocal brain damage induced by phencyclidine is augmented by pilocarpine. *Brain Res* 33:533–541.
- DeOlmos JS, Ingram WR (1971) An improved cupric-silver method for impregnation of axonal and terminal degeneration. *Brain Res* 33:523–529.
- Dikranian K, Ishimaru MJ, Tenkova T, Labruyere J, Qin YQ, Ikonomidou C, Olney JW (2001) Apoptosis in the in vivo mammalian forebrain. *Neurobiol Dis* 8:359–379.
- Dobbing J, Sands J (1979) The brain growth spurt in various mammalian species. *Early Hum Dev* 3:79–84.
- Franks NP, Lieb WR (1994) Molecular and cellular mechanism of general anesthesia. *Nature* 367:607–614.
- Franks NP, Dickinson R, deSousa SLM, Hall AC, Lieb WR (1998) How does xenon produce anesthesia? *Nature* 396:324.
- Gonsowski CT, Eger II EI (1994) Nitrous oxide minimum alveolar anesthetic concentration in rats is greater than previously reported. *Anesth Analg* 79:710–712.
- Hartman RE, Wozniak DF, Nardi A, Olney JW, Sartorius L, Holtzman DM (2001) (2002) Behavioral phenotyping of GFAP-*apoE3* and -*apoE4* transgenic mice: *ApoE4* mice show profound working memory impairments in the absence of Alzheimer's-like neuropathology. *Exp Neurol* 170:326–344.
- Ho N, Liauw JA, Blaeser F, Wei F, Hanissian S, Muglia LM, Wozniak DF, Nardi A, Linden DJ, Zhuo M, Muglia LJ, Chatila TA (2000) Impaired synaptic plasticity and CREB activation in *CaMKIV/Gr*-deficient mice. *J Neurosci* 20:6459–6472.
- Hornbein TF, Eger II EI, Winter PM, Smith G, Wetstone D, Smith KH (1982) The minimum alveolar concentration of nitrous oxide in man. *Anesth Analg* 61:553–556.
- Ikonomidou C, Bosch F, Miksa M, Bittigau P, Vockler J, Dikranian K, Tenkova TI, Stefovskaya V, Turski L, Olney JW (1999) Blockade of NMDA receptors and apoptotic neurodegeneration in the developing brain. *Science* 283:70–74.
- Ikonomidou C, Bittigau P, Ishimaru MJ, Wozniak DF, Koch C, Genz K, Price MT, Stefovskaya V, Horster F, Tenkova T, Dikranian K, Olney JW (2000) Ethanol-induced apoptotic neurodegeneration and fetal alcohol syndrome. *Science* 287:1056–1060.
- Ishimaru MJ, Ikonomidou C, Tenkova TI, Der TC, Dikranian K, Sesma MA, Olney JW (1999) Distinguishing excitotoxic from apoptotic neurodegeneration in the developing rat brain. *J Comp Neurol* 408:461–476.
- Jevtovic-Todorovic V, Todorovic SM, Mennerick S, Powell S, Dikranian K, Benshoff N, Zorumski CF, Olney JW (1998) Nitrous oxide (laughing

- gas) is an NMDA antagonist, neuroprotectant, and neurotoxin. *Nat Med* 4:460–463.
- Kissin I, Brown PT, Bradley EL (1990) Sedative and hypnotic midazolam-morphine interactions in rats. *Anesth Analg* 71:137–143.
- Lodge D, Anis NA (1982) Effects of phencyclidine on excitatory amino acid activation of spinal interneurons in the cat. *Eur J Pharmacol* 77:203–204.
- Mahmoudi NW, Cole DJ, Shapiro HM (1989) Insufficient anesthetic potency of nitrous oxide in the rat. *Anesthesiology* 70:345–349.
- Mennerick S, Jevtovic-Todorovic V, Todorovic SM, Shen W, Olney JW, Zorumski CF (1998) Effect of nitrous oxide on excitatory and inhibitory synaptic transmission in hippocampal cultures. *J Neurosci* 18:9716–9726.
- Mitchell AS, Dalrymple-Alford JC, Christie MA (2002) Spatial working memory and the brainstem cholinergic innervation to the anterior thalamus. *J Neurosci* 22:1922–1928.
- Olney JW, Tenkova TI, Dikranian K, Muglia LJ, Jermakowicz WJ, D'Sa C, Roth KA (2002a) Ethanol-induced apoptotic neurodegeneration in the developing C57BL/6 mouse brain. *Brain Res Dev Brain Res* 133:115–126.
- Olney JW, Tenkova TI, Dikranian K, Qin YQ, Labruyere J, Ikonomidou C (2002b) Ethanol-induced caspase-3 activation in the in vivo developing mouse brain. *Neurobiol Dis* 9:205–219.
- Orliaguet G, Vivien B, Langeron O, Bouhemad B, Coriat P, Riou B (2001) Minimum alveolar concentration of volatile anesthetics in rats during postnatal maturation. *Anesthesiology* 95:734–739.
- Stevens WC, Kingston HGG (1992) Inhalation anesthesia. In: *Clinical anesthesia* (Barash PG, Cullen BF, Stoelting RK, eds), pp 385–412. Philadelphia: Lippincott.
- Streissguth AP, O'Malley K (2000) Neuropsychiatric implications and long-term consequences of fetal alcohol spectrum disorders. *Semin Clin Neuropsychiatry* 5:177–190.
- West MJ (1999) Stereological methods for estimating the total number of neurons and synapses: issues of precision and bias. *Trends Neurosci* 22:51–61.
- Wozniak DF, Stewart GR, Finger S, Olney JW (1989) Comparison of behavioral effects of nucleus basalis magnocellularis lesions and somatosensory cortex ablation in the rat. *Neuroscience* 32:685–700.
- Wozniak DF, Olney JW, Kettinger III L, Price M, Miller JP (1990) Behavioral effects of MK-801 in the rat. *Psychopharmacology* 101:47–56.
- Wozniak DF, Cicero TJ, Kettinger III L, Meyer ER (1991) Paternal alcohol consumption in the rat impairs spatial learning performance in male offspring. *Psychopharmacology* 105:289–302.
- Zorumski CF, Mennerick S, Izumi Y (1996) Assessment of synaptic effects of nitric oxide in hippocampal neurons. *Methods Neurosci* 31:283–299.